

BRIEF REPORT

Simulating Murder: The Aversion to Harmful Action

Fiery Cushman
Harvard University

Kurt Gray
The University of Maryland

Allison Gaffey
University of Notre Dame

Wendy Berry Mendes
University of California, San Francisco

Diverse lines of evidence point to a basic human aversion to physically harming others. First, we demonstrate that unwillingness to endorse harm in a moral dilemma is predicted by individual differences in aversive reactivity, as indexed by peripheral vasoconstriction. Next, we tested the specific factors that elicit the aversive response to harm. Participants performed actions such as discharging a fake gun into the face of the experimenter, fully informed that the actions were pretend and harmless. These simulated harmful actions increased peripheral vasoconstriction significantly more than did witnessing pretend harmful actions or to performing metabolically matched nonharmful actions. This suggests that the aversion to harmful actions extends beyond empathic concern for victim harm. Together, these studies demonstrate a link between the body and moral decision-making processes.

Keywords: morality, aggression, aversion, empathy, psychophysiology

People are averse to performing harmful actions and often consider it morally wrong to harm a person even when it would save many more lives (Mikhail, 2000; Petrinovich, O'Neill, & Jorgensen, 1993). Even frontline soldiers trained and motivated to kill often deliberately miss visible enemy targets (Grossman, 1995). This aversion to harm is essential to ordinary human functioning, as evidenced by the antisocial behavior of psychopaths, who are argued to lack it (Blair, 1995). The aversion to harming others is so basic to our moral sense that it is easy to miss an important question: What is its psychological basis?

Our first experiment examines the link between physiological responses and answers on a classic moral dilemma: whether killing someone is allowable to save many lives. We measured physiological reactivity, linked to general aversive states, during a non-moral task and then examined whether it predicted advocating the

death of one person to save others. We used autonomic changes, specifically changes in total peripheral resistance (TPR), which are associated with negative stress responses (Gregg, James, Matyas, & Thorsteinsson, 1999; Mendes, Blascovich, Hunter, Lickel, & Jost, 2007). Past research indicates that the aversion to harm in such moral dilemmas involves an affective component (e.g., Greene, Sommerville, Nystrom, Darley, & Cohen, 2001; Mendez et al., 2005; Moretto, Ládavas, Mattioli, & di Pellegrino, 2010) but does not establish a link between the specific aversion to harm in moral dilemmas and general aversive reactivity in nonmoral situations.

Our second experiment builds on this finding, asking why people find the performance of harmful actions aversive. First, aversion may stem from empathic concern for the welfare of the victim (Crockett, Clark, Hauser, & Robbins, 2010; Hoffman, 2000; Mehrabian & Epstein, 1972; Pizarro, 2000). For instance, we might be averse to punching another person because considering the victim's pain causes us psychological distress (Batson et al., 2003; Gray, Gray, & Wegner, 2007; Singer et al., 2004). It is important to note that aversion is not intrinsic to the action itself but rather to its expected outcome. We call this the *outcome aversion* model: People are averse to harmful acts because of empathic concern for victim distress.

In addition, an aversive response might be triggered by the basic perceptual and motoric properties of an action, even without considering its outcome. Blair (1995) has suggested a mechanism by which harmful actions themselves can become aversive: When the unconditioned aversive stimulus of victim distress (e.g., crying) is repeatedly paired with a particular action (e.g., pushing or hitting a person), those actions acquire a conditioned aversive response. On this *action aversion* model, empathy is critical to the acquisi-

This article was published Online First September 12, 2011.

Fiery Cushman, Department of Psychology, Harvard University; Kurt Gray, Department of Psychology, The University of Maryland; Allison Gaffey, Department of Psychology, University of Notre Dame; and Wendy Berry Mendes, Department of Psychiatry, University of California, San Francisco.

Fiery Cushman is now at Brown University.

We thank our undergraduate research assistants and other members of the Health and Physiology Laboratory and Moral Cognition Laboratory for their assistance in collecting the data. This research was supported by Grant HL079383 from the National Heart, Lung, and Blood Institute and funding from the Mind, Brain and Behavior Initiative at Harvard University.

Correspondence concerning this article should be addressed to Fiery Cushman, Department of Cognitive, Linguistic and Psychological Sciences, Box 1821, Brown University, Providence, RI 02912. E-mail: fiery_cushman@brown.edu

tion of the aversive response to harmful actions, but the conditioned response may subsequently be evoked by intrinsic properties of the action alone.

In Study 2, we tested for action aversion by examining participants' physiological responses while either performing or witnessing harmful actions (stabbing an experimenter with a rubber knife, shooting him with a disabled handgun, etc.) or performing similar but harmless actions (e.g., slicing a pretend loaf of bread with a knife). Our use of simulated actions follows past research demonstrating that pretend stimuli can be sufficient to elicit strong psychological responses (Rozin, Millman, & Nemeroff, 1986). Action aversion predicts a robust aversive response to pretend actions with motoric and perceptual properties of actual harmful behaviors although the "perpetrator" knows that no harm will occur, whereas outcome aversion does not. Additionally, action aversion predicts a greater aversive response to performing harm than witnessing it (because only the former involves an action), whereas outcome aversion predicts an equal aversive response in both cases (because they yield the same outcome).

In summary, in Study 1, we tested the relationship between the moral judgment of harmful actions and general TPR reactivity to a nonmoral task. In Study 2, we tested whether simulated harms specifically trigger TPR reactivity. Additionally, we compared reactivity for performing versus witnessing simulated harm, testing whether reactivity depends on the anticipation of a harmful outcome versus the performance of a harmful act.

Study 1

Study 1 examines the relationship between threat reactivity and responses to a classic moral dilemma. We tracked changes in TPR during a stressful arithmetic task and predicted that individuals exhibiting greater TPR reactivity would be less willing to endorse harming one person to save the lives of several others.

Method

We recruited 108 healthy participants (81 female, 27 male) aged 19–40 years ($Mdn = 24$). After obtaining consent, an experimenter applied sensors that measured impedance cardiography (HIC 2500, Chapel Hill, NC), electrocardiography (Biopac ECG module, Goleta, CA), and blood pressure responses (Colin Prodigy II, San Antonio, TX). Impedance cardiographic and electrocardiography signals were sampled at 1000 Hz and integrated with a Biopac MP150. Postacquisition waveforms were scored with Mindware software (IMP 3.0) by trained research assistants (see Mendes, 2009). We estimated TPR using the standard formula:

$$TPR = (\text{mean arterial pressure} / \text{cardiac output}) \times 80.$$

After baseline, participants met a new experimenter who asked them to count backward quickly in steps of 7 from a four-digit number. Mental arithmetic is a common laboratory stress task that can evoke increases in sympathetic nervous system responding. We used TPR change during the first minute of the stress task as our indication of threat reactivity, subtracting the last minute of the baseline period from the first minute of the stress task.

Participants were recruited as part of a larger study on physiological changes associated with emotion and body manipulations, analyses of which are beyond the scope of the present study. Here

we report physiological responses that occur before our emotion manipulations. The body position manipulation (leaning forward vs. leaning back) was introduced before the stress task but had no effect on the early physiological responses we analyze here.

Participants were then provided a packet of questionnaires that included a moral dilemma. It asked the participant to imagine being on a lifeboat that would sink—killing all onboard—unless someone is thrown off. One person on the lifeboat is "leaning over the side." Participants were asked, "Is it morally acceptable for you to push this person overboard in order to save the lives of the remaining passengers?" indicating a yes-or-no answer. Then they were asked to indicate "how morally acceptable it would be for you to throw this person overboard in order to save the lives of the remaining passengers" on a scale ranging from 1 (*completely unacceptable*) to 7 (*completely acceptable*).

Results and Discussion

Measurement of TPR requires several high-quality, artifact-free physiological signals. Twenty participants were excluded for low-quality impedance or electrocardiograph waveform; 23, because blood pressure measurements were not obtained during the first minute of the task; 3, because TPR reactivity scores differed by more than 2 standard deviations from the mean; and another 9 dropped out of the study before the assessment of the moral dilemmas. Responses to moral dilemmas did not significantly differ between those with usable TPR data and those without, $t(106) = 0.60, ns$.

As predicted, increased TPR reactivity was reliably associated with lesser endorsement of pushing a person overboard in our moral judgment task $r = -.31, n = 51, p < .05$, although not when rated dichotomously, $t(40) < .01, ns$. This result was robust after controlling for the experimental manipulations of posture, affect, their interaction, gender, and age, $r = -.31, n = 51, p < .05$.¹

The observed correlation between moral judgment and TPR reactivity is consistent with our prediction that unwillingness to endorse harmful action is linked with threat reactivity. This suggests that the aversion to harmful actions may be instantiated physiologically. However, it leaves open the basis of this response: Does it depend solely on empathy for an actual victim or also on the perceptual and motoric properties of the action itself?

Study 2

Study 2 tested the action aversion hypothesis—whether physiological aversion can be triggered by only the motor or perceptual properties of harmful action. Participants were asked to perform five simulated harmful actions, to witness another person perform them, or to perform five simulated nonharmful actions. We tested two predictions of the action aversion hypothesis: Simulated harmful actions would elicit aversive reactivity despite the absence of

¹ We also examined the association between responses on moral dilemmas and changes in cardiac output. Consistent with the "threat" profile, cardiac output decreases were associated with less endorsement of harming others, $r = .29, n = 54, p < .04$. In Study 2, we examine responses during a task that does not meet the requirements of a motivated performance situation, so we focus on TPR in both studies for consistency.

any harmful outcome, and this response would be greater when performing the action than when witnessing the action.

Method

We recruited 108 participants (69 female, 39 male) aged 18–35 years ($Mdn = 20$). Participants initially consented to a study of “pretend actions” omitting any mention of harm. Participants completed 20 items from the Positive and Negative Affect Schedule (PANAS; Watson, Clark, & Tellegen, 1988), and then we applied sensors from two different ambulatory devices that allowed free movement during testing: VU-AMS impedance/ECG (Amsterdam), and SpaceLab blood pressure (ABP 90207, Issaquah, WA). The experimenter measured baseline physiological responses during a 5-min rest and then brought participants into a room where she described the full experimental procedure, obtained additional informed consent which emphasized that the participant was free to omit any actions, and then initiated the pretend actions.

A male research assistant played the role of victim in the perform-harm and witness-harm conditions. Participants in the perform-harm condition were asked to perform five actions in a fixed order: (1) smashing the victim’s shin with a hammer—a PVC pipe was worn under a fake pant leg, (2) smashing the victim’s hand with a rock—a rubber hand was placed at the cuff of the shirt and the actual hand was obscured from sight, (3) discharging a handgun into the victim’s face—a weighty metal replica, (4) drawing a knife across the victim’s throat—a rubber knife, and (5) smacking a baby against the table—we used a realistic looking baby doll (see Figure 1). No verbal communication occurred between the participant and the victim, and the victim avoided eye contact except during the action itself. The victim grimaced slightly during each action but exhibited no further distress.

The experimenter initiated each action by describing it to the participant and emphasizing that it was pretend and that no harm

would occur. The participant was instructed to contemplate performing the action for 1 min while holding the relevant implement (e.g., gun), during which time the experimenter exited the room. The experimenter returned and said, “It is time to perform the action.” After the participant performed (or chose not to perform) the action, the experimenter closed a curtain between the participant and the victim, instructed the participant to sit quietly for 1 min, and exited. This sequence was repeated for each action.

The witness-harm condition proceeded identically, except that the participant was introduced to two additional experimenters who assumed the roles of perpetrator and victim. The participant heard an identical description of the event to be performed and then contemplated watching that event for 1 min. The experimenter returned to the room and asked the perpetrator to “harm” the victim, which the perpetrator did directly in front of the participant with neutral affect. The perpetrator and the victim were then masked by a curtain during a 1-min postaction period. The no-harm condition also proceeded identically, except that there were no additional experimenters, and the participant was asked to perform five metaphorically controlled pretend actions: (1) hammering an imaginary nail on a block of wood, (2) using a rock to smash a (rubber) nut, (3) using a spray bottle to mist an imaginary plant, (4) using a rubber knife to cut a (cardboard) loaf of bread, and (5) smacking a hand broom against a table to shake out dust.

After the pretend actions, participants returned to the original room and sensors were removed. Participants completed several questionnaires including the 20-item PANAS and five hypothetical moral dilemmas drawn from previous research (Greene et al., 2001). These moral dilemmas asked participants whether they would perform a harmful action to save many lives (e.g., whether to smother one’s own crying baby to successfully hide their whole family from enemy soldiers).




Figure 1. Harmful and nonharmful actions used in Experiment 2.

During the experiment, impedance and electrocardiograph were monitored continuously, but blood pressure measurements were manually initiated by the experimenter with a key press. We took blood pressure readings at the first, third, and fifth actions, and the timing of these (before vs. after the action) was varied between participants. Thus, for each participant, we calculated three TPR reactivity scores.

One participant elected not to perform any actions and was excluded from all analyses. Thirty-three participants were excluded from physiological analyses because of equipment malfunction, experimenter error, or biologically implausible measurements. Individual data points greater than 2 standard deviations from the group mean were excluded from analysis. Several participants were excluded from behavioral analysis because of incomplete responses to posttest survey items.

Results and Discussion

Self-reported affect. An analysis of covariance of posttask negative affect (controlling for pretask scores) yielded a significant effect, $F(2, 101) = 16.90, p < .001$. Simple effects revealed increased negative affect for perform-harm ($p < .001$) and witness-harm ($p < .001$) conditions compared with the no-harm condition but no difference between the perform and witness conditions ($p > .25$).

Harmful actions. We used the general linear model to test for effects of condition (perform harm vs. witness harm vs. no harm) and measurement period (preaction contemplation vs. postaction recovery) on TPR reactivity, treating participant as a random effect and using robust standard errors. There was a significant effect of measurement period, $\beta_1 = -0.31, z = -3.30, p < .01$: Across all three conditions, TPR reactivity was greater preaction than it was postaction. We treated the perform-harm condition as the comparison condition and found that TPR reactivity was significantly greater in this condition than it was in the witness-harm condition, $\beta_2 = -0.23, z = -2.08, p < .05$; and the no-harm condition, $\beta_3 = -0.26, z = -2.65, p < .01$ (see Table 1). Additional analyses revealed no significant effect of order (earlier vs. later actions) on TPR reactivity and no significant interactions between condition, measurement period, or task order.

We performed a supplementary analysis of TPR reactivity taken from the very first preaction contemplation period, a point at which participants had anticipated the task but had not performed or witnessed any action. TPR reactivity for the perform-harm condition ($M = 89.8$) was significantly higher than it was for the

witness-harm condition ($M = 30.7$), $\beta = -0.37, t = 2.36, p < .05$; and for the no-harm condition ($M = 29.9$), $\beta = -0.37, t = 2.34, p < .05$.

Moral dilemmas. Paralleling our analysis in Study 1, we assessed whether greater TPR reactivity was associated with lesser endorsement of harming one person to save several others. We calculated a summary TPR score for each participant, averaging their reactivity measurements and adjusting postaction TPR scores to match preaction TPR scores according to the relevant coefficient (β_1) of the generalized linear model presented earlier. We then correlated TPR reactivity with the mean judgment across five moral dilemmas. Collapsed across all three conditions, this correlation was significant, $r = -.32, n = 71, p < .01$. The relationship was larger and significant for the witness-harm condition, $r = -.49$ (95% confidence interval [CI] = $-.12, -.87$), $n = 25, p < .05$; smaller and nonsignificant for the perform-harm condition, $r = -.36$ (95% CI = $-.06, .78$), $n = 23, p < .10$; and smallest and nonsignificant for the no-harm condition, $r = -.24$ (95% CI = $-.20, .68$), $n = 23, p < .28$. An analysis of variance revealed no significant effect of condition on mean moral judgment, $F(2, 97) = 0.75, p = .47$. Although our condition effects did not influence later moral judgments on hypothetical scenarios, individual differences in TPR reactivity predicted moral judgments as in Study 1.

General Discussion

We investigated individuals' aversion to harmful actions. Study 1 demonstrated that individuals exhibiting greater threat reactivity were less likely to endorse harm to save lives. This finding corroborates past evidence suggesting a more potent aversive response to the idea of performing direct harm than to allowing indirect harm to more distant others. It also suggests that individual differences in TPR during nonstressful tasks can predict this aversion to harm.

Study 2 investigated the psychological basis of this aversive response. Performing simulated harmful actions evoked robust TPR reactivity despite the participants' full awareness that no actual harm would be caused. TPR reactivity was lower among participants who were asked to witness harmful actions or to perform metabolically controlled nonharmful actions. Moreover, TPR reactivity differed between conditions during the very first preaction measurement period, before participants performed or witnessed any action at all. Thus, simply contemplating performing a simulated harmful action leads to greater vasoconstriction than contemplating witnessing a harmful action.

These findings suggest an aversion to performing harmful actions that extends beyond the expectation of a harmful outcome. Clearly, TPR reactivity to pretend harmful action (e.g., hitting a plastic baby doll) cannot be attributed to the explicit belief that harm will occur. Nevertheless, pretend events could trigger the imagination of harmful outcomes. Critically, however, outcome aversion predicts similar affective states for witnessing and performing harm, whereas action aversion predicts a unique aversive response to performing harm, as we observed. We therefore consider it unlikely that the TPR reactivity associated with performing simulated harmful actions was caused solely by consideration of a harmful outcome, such as empathic concern for victim distress.

Table 1
Mean Total Peripheral Resistance (TPR) Reactivity by
Condition and Measurement Period

Condition	TPR change: M (SD, n)	
	Preaction	Postaction
Perform harm	75 (73, 34)	47 (90, 28)
Witness harm	34 (73, 49)	10 (88, 24)
No harm	29 (83, 48)	-28 (68, 16)

Note. n = number of observations, with up to three observations per participant.

Our findings do not contradict the role of empathy and victim distress on the aversion to harmful actions. These elements may be especially important for the performance of “real” harmful actions with actual consequences; moreover, they may play a key developmental acquisition of action aversion through associative pairing (Blair, 1995). Just as you cannot help but swoon when smelling the perfume or cologne associated with your first love, people cannot help but feel upset when doing actions typically associated with victim distress. However, as important as the aversion to victim distress may be, our results suggest a dissociable aversion based on mere actions.

A forceful, automatic aversive response to the surface properties of harmful actions may explain otherwise puzzling human behaviors. In battlefield behavior and hypothetical moral judgment, people resist doing direct harm despite explicit knowledge that it could save many lives. Similarly, in our study, people experienced a strong aversive response to performing pretend harmful actions despite the explicit knowledge that no harm would be caused. These cases highlight a dissociation between our explicit knowledge of the consequences of our actions and our automatic affective responses to actions (Dayan & Niv, 2008; Kahneman, 2003; LeDoux, 1996).

The action aversion model also suggests a darker side: When banal or novel actions lack motoric and perceptual properties associated with harm, they may fail to trigger an aversive response. Signing one’s name to a torture order or pressing the button that releases a bomb each have real, known consequences for other people, but as actions they lack salient properties reliably associated with victim distress. A notable parallel is evident in moral judgment: People consider it morally worse to cause harm through direct physical engagement than at a distance (Cushman, Young, & Hauser, 2006; Greene et al., 2009). We demonstrate that TPR reactivity increases during (pretend) harmful actions and also correlates with judgments of moral dilemmas. However, although circumstantial evidence implicates a role for action aversion in moral judgment, further research is required.

As such, our study highlights the advantage of taking lessons from hypothetical moral dilemmas and translating them into more active behaviors. Few studies have directly targeted the human aversion to harm using an active behavioral paradigm (Martens, Kosloff, Greenberg, Landau, & Schmader, 2007; Milgram, 1974). This is no surprise: It is hard to get one person to harm another ethically and in a laboratory. Moreover, past studies often targeted situational factors that promote harm, rather than the affective systems that discourage it. To ask why people *do* harm is a critical research question; our complementary question is why people *do not*. Our study suggests that the use of simulated harmful actions is sufficient to generate an aversive response. Surely this response is weaker than the aversion experienced by a soldier on the battlefield or the captain of a sinking ship; nevertheless, the aversion to simulated harm in the laboratory may provide insight into the psychology underlying the aversion to actual harm in the world beyond.

References

- Batson, D. C., Lishner, D. A., Carpenter, A., Dulin, L., Harjusola-Webb, S., Stocks, E. L., . . . Sapat, B. (2003). “As you would have them do onto you”: Does imagining yourself in the other’s place stimulate moral action? *Personality and Social Psychology Bulletin*, *29*, 1190–1201. doi:10.1177/0146167203254600
- Blair, R. J. R. (1995). A cognitive developmental approach to morality: Investigating the psychopath. *Cognition*, *57*, 1–29. doi:10.1016/0010-0277(95)00676-P
- Crockett, M., Clark, L., Hauser, M., & Robbins, T. (2010). Serotonin selectively influences moral judgment and behavior through effects on harm aversion. *Proceedings of the National Academy of Sciences, USA*, *107*, 17433. doi:10.1073/pnas.1009396107
- Cushman, F. A., Young, L., & Hauser, M. D. (2006). The role of conscious reasoning and intuitions in moral judgment: Testing three principles of harm. *Psychological Science*, *17*, 1082–1089. doi:10.1111/j.1467-9280.2006.01834.x
- Dayan, P., & Niv, Y. (2008). Reinforcement learning: The good, the bad and the ugly. *Current Opinion in Neurobiology*, *18*, 185–196. doi: 10.1016/j.conb.2008.08.003
- Gray, H. M., Gray, K., & Wegner, D. M. (2007). Dimensions of mind perception. *Science*, *315*, 619. doi:10.1126/science.1134475
- Greene, J. D., Cushman, F. A., Stewart, L. E., Lowenberg, K., Nystrom, L. E., & Cohen, J. D. (2009). Pushing moral buttons: The interaction between personal force and intention in moral judgment. *Cognition*, *111*, 364–371. doi:10.1016/j.cognition.2009.02.001
- Greene, J. D., Sommerville, R. B., Nystrom, L. E., Darley, J. M., & Cohen, J. D. (2001). An fMRI investigation of emotional engagement in moral judgment. *Science*, *293*, 2105–2108. doi:10.1126/science.1062872
- Gregg, M. E., James, J. E., Matyas, T. A., & Thorsteinsson, E. B. (1999). Hemodynamic profile of stress-induced anticipation and recovery. *International Journal of Psychophysiology*, *34*, 147–162. doi:10.1016/S0167-8760(99)00074-4
- Grossman, D. (1995). *On killing*. Boston: Little, Brown.
- Hoffman, M. L. (2000). *Empathy and moral development*. Cambridge, United Kingdom: Cambridge University Press.
- Kahneman, D. (2003). Maps of bounded rationality: Psychology for behavioral economics. *American Economic Review*, *93*, 1449–1475. doi: 10.1257/00028280322655392
- LeDoux, J. (1996). *The emotional brain*. New York: Simon and Shuster.
- Martens, A., Kosloff, S., Greenberg, J., Landau, M., & Schmader, T. (2007). Killing begets killing: Evidence from a bug-killing paradigm that initial killing fuels subsequent killing. *Personality and Social Psychology Bulletin*, *33*, 1251.
- Mehrabian, A., & Epstein, N. (1972). A measure of emotional empathy. *Journal of Personality*, *40*, 525–543. doi:10.1111/j.1467-6494.1972.tb00078.x
- Mendez, M. F., Anderson, E., & Shapria, J. S. (2005). An investigation of moral judgment in frontotemporal dementia. *Cognitive and Behavioral Neurology*, *18*, 193–197.
- Mendes, W., Blascovich, J., Hunter, S., Lickel, B., & Jost, J. (2007). Threatened by the unexpected: Physiological responses during social interactions with expectancy-violating partners. *Journal of Personality and Social Psychology*, *92*, 698. doi:10.1037/0022-3514.92.4.698
- Mendes, W. B. (2009). Assessing the autonomic nervous system. In: E. Harmon-Jones and J. Beer (Eds.) *Methods in the Neurobiology of Social and Personality Psychology*. Guilford Press.
- Mikhail, J. M. (2000). *Rawls’ linguistic analogy: A study of the “generative grammar” model of moral theory described by John Rawls in “A Theory of Justice.”* Unpublished doctoral dissertation, Cornell University, Ithaca, NY.
- Milgram, S. (1974). *Obedience to authority: An experimental view*. New York: Harper & Row.
- Moretto, G., Làdavas, E., Mattioli, F., & di Pellegrino, G. (2010). A psychophysiological investigation of moral judgment after ventromedial prefrontal damage. *Journal of Cognitive Neuroscience*, *22*, 1888–1899. doi:10.1162/jocn.2009.21367
- Petrinovich, L., O’Neill, P., & Jorgensen, M. J. (1993). An empirical study of moral intuitions: Towards an evolutionary ethics. *Journal of Personality and Social Psychology*, *64*, 467–478. doi:10.1037/0022-3514.64.3.467

- Pizarro, D. A. (2000). Nothing more than feelings? The role of emotions in moral judgment. *Journal for the Theory of Social Behavior*, 30, 355–375. doi:10.1111/1468-5914.00135
- Rozin, P., Millman, L., & Nemeroff, C. (1986). Operation of the laws of sympathetic magic in disgust and other domains. *Journal of Personality and Social Psychology*, 50, 703–712. doi:10.1037/0022-3514.50.4.703
- Singer, T., Seymour, B., O’Doherty, J., Kaube, H., Dolan, R. J., & Frith, C. D. (2004). Empathy for pain involves the affective but not sensory components of pain. *Science*, 303, 1157–1162.
- Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*, 54, 1063–1070. doi:10.1037/0022-3514.54.6.1063

Received February 11, 2011

Revision received April 4, 2011

Accepted April 5, 2011 ■

Members of Underrepresented Groups: Reviewers for Journal Manuscripts Wanted

If you are interested in reviewing manuscripts for APA journals, the APA Publications and Communications Board would like to invite your participation. Manuscript reviewers are vital to the publications process. As a reviewer, you would gain valuable experience in publishing. The P&C Board is particularly interested in encouraging members of underrepresented groups to participate more in this process.

If you are interested in reviewing manuscripts, please write APA Journals at Reviewers@apa.org. Please note the following important points:

- To be selected as a reviewer, you must have published articles in peer-reviewed journals. The experience of publishing provides a reviewer with the basis for preparing a thorough, objective review.
- To be selected, it is critical to be a regular reader of the five to six empirical journals that are most central to the area or journal for which you would like to review. Current knowledge of recently published research provides a reviewer with the knowledge base to evaluate a new submission within the context of existing research.
- To select the appropriate reviewers for each manuscript, the editor needs detailed information. Please include with your letter your vita. In the letter, please identify which APA journal(s) you are interested in, and describe your area of expertise. Be as specific as possible. For example, “social psychology” is not sufficient—you would need to specify “social cognition” or “attitude change” as well.
- Reviewing a manuscript takes time (1–4 hours per manuscript reviewed). If you are selected to review a manuscript, be prepared to invest the necessary time to evaluate the manuscript thoroughly.